

Routine Schedule

Age*	Vaccine	Primary Doses	Booster Dose
6 months– 4 years	Pfizer – Infant/Toddler	→ 3-8 weeks [^] → → ≥8 weeks →	
6 months– 5 years	Moderna – Infant/Toddler	→ 4-8 weeks [^] →	
5–11 years	Pfizer – Pediatric	→ 3-8 weeks [^] → → ≥5 months →	
6–11 years	Moderna – Pediatric	→ 4-8 weeks [^] →	
12+ years	Moderna – Adol/Adult	→ 4-8 weeks [^] →	
12+ years	Novavax	→ 3-8 weeks [^] →	
12+ years	Pfizer/ Adol/Adult	→ 3-8 weeks [^] →	
18+ years	Janssen (J&J) Pfizer/Moderna preferred**		
























* See schedules for children in transition from a younger to older age group: [Pfizer](#) | [Moderna](#).

** Although use of mRNA COVID-19 vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

[^] An [8-week interval](#) may be preferable for some people, especially for males 12-39 years.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

Schedule if Moderately or Severely Immunocompromised

Age*	Vaccine	Primary Doses	Booster Dose
6 months– 4 years	Pfizer– Infant/Toddler	 3 weeks  ≥8 weeks 	
6 months– 5 years	Moderna– Infant/Toddler	 4 weeks  ≥4 weeks 	
5–11 years	Pfizer– Pediatric	 3 weeks  ≥4 weeks 	 <div style="border: 1px solid black; padding: 5px; background-color: #e0f2f1;"> Monovalent Booster Pfizer 5-11 years </div>
6–11 years	Moderna– Pediatric	 4 weeks  ≥4 weeks 	
12+ years	Moderna– Adol/Adult	 4 weeks  ≥4 weeks 	 <div style="border: 1px solid black; padding: 10px; background-color: #e0f2f1;"> Bivalent Booster Pfizer: Ages 12+ Moderna: Ages 18+ (Regardless of the number of monovalent booster doses previously received) </div>
12+ years	Novavax	 3 weeks 	
12+ years	Pfizer/ Adol/Adult	 3 weeks  ≥4 weeks 	
18+ years	Janssen (J&J) Pfizer/Moderna preferred**	 4 weeks <div style="border: 1px solid black; padding: 5px; background-color: #e0f2f1; display: inline-block;"> 2nd Dose of Moderna/ Pfizer </div>	

* See schedules for children in transition from a younger to older age group: [Pfizer](#) | [Moderna](#).

** Although use of mRNA COVID-19 vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.